

A Review on Tannase and its Applications

Archana Tripathi¹, B Lakshmi²

How to cite this article:

Archana Tripathi, B Lakshmi/ A Review on Tannase and its Applications/ J Microbiol Relat Res. 2021;7(1): 29–39.

Abstract

Tannin acyl hydrolase (E.C. 3.1.1.20) or Tannase is a type of hydrolase enzyme that catalyzes the hydrolysis of tannin and is converted to gallic acid and glucose. Natural tannins occur in various parts of plants such as leaves, roots, bark and fruits. Tannins are distributed in animals, insects and microorganisms. However, microbial tannase is one of the great attention in many industries. Gallic acid is one of the industrial and therapeutic molecules that is widely used in various applications. Tannase is also applicable in feed, pharmaceutical, brewing, chemical and beverage industries. Furthermore, tannase plays a significantly important role in biodegradation or reduction of tannin from the tannery effluent. The present review appraises a brief description of tannase and its industrial applications.

Keywords: Tannin; Tannase; Gallic acid; Glucose; Tannery effluent.

Introduction

Tannases represent a group of enzymes, that are produced by microbes, plants and animals. However, tannase from microbial sources is preferred over other sources for industrial. The catalytic reaction of tannase involves the hydrolysis of ester bond and depside bond present in various substrates i.e., tannic acid, complex tannin, gallic acid esters, epigallocatechin gallate, methyl gallate, ethyl gallate, n-propyl gallate to produce gallic acid (3, 4, 5-tri hydroxybenzoic acid) and glucose. Various parameters such as temperature, pH, metal ions, activators, inhibitors and substrate specificity

affect the enzyme activity.

Tannase enzymes are used in the food, brewing, and pharmaceutical industries. Gallic acid production is one of the most important commercial applications of tannase. Apart from that, they are extensively used in the food industry, especially in instant tea production, where it enhances the extractability and cold water solubility of key compounds. Another important application of tannase is the removal of haze formation and unflavored phenolic compounds from beer and wine and fruit juices. The quality of fruit juices also can be improved by the tannase enzyme.

Industries require novel microbial strains for enzyme production using easily available inexpensive raw materials and trying to evolve various dimensions of tannase such as novel sources of tannase, new microbes with high tannin transformation ability, new substrate specificities, low cost production, less expenditure on the purification strategies and the potential use of tannase in different industries has been shown by researchers through out the world. The current review articles explain in detail the structure, purification and application of tannase produced by microbes.

Author Affiliation: ¹Lecturer, ²Assistant Professor, Department of Biotechnology and Microbiology, Shri Maneklal M Patel Institute of Science and Research, Kadi Sarva Vishwavidyalaya, Gandhinagar 382015 Gujarat, India.

Corresponding Author: B Lakshmi, Assistant Professor, Department of Biotechnology and Microbiology, Shri Maneklal M Patel Institute of Science and Research, Kadi Sarva Vishwavidyalaya, Gandhinagar 382015 Gujarat, India.

E-mail: lakshmi.smmprsr@ksv.ac.in

Received on: 12-03-2022

Accepted on: 21.03.2022

Tannins

The meaning of Tannin or Tanna (German word) is oak, which refers to the use of tannins from the wood/bark of oak or fir trees that were used in the tanning process (Barbehenn and Constabel, 2011; Tomak and Gonultas, 2018). Tannins are well known large, complex, natural plant derived polyphenolic biomolecules that are found in the bark, stem, buds, wood, leaves, fruits and roots. Tannin plays a very important role in protecting plants from microorganisms and other predators (Khanbabaee and van Ree, 2001; Sharma, 2019). In a few countries, tannin rich plant extracts serve as traditional medicines for the treatment of cancer, diarrhea, inflammation, diuresis, hemorrhagic stroke and skin (Dai, 2010 and Działo, 2016).

Based on diversified structures and properties of tannins, they are classified into three major classes and two subclasses: (1) Hydrolysable tannins (2) Condensed tannins and (3) Complex tannins (Figure 1). Hydrolysable tannins can be defined as the types of tannins that are hydrolysed into

gallic acid and ellagic acid in the presence of acid or alkaline conditions, hot water and enzymes. Hydrolysable tannins are further classified into two subclasses based on their hydrolysable products i.e. (i) Gallotannins and (ii) Ellagitannins. Hydrolysable tannins are mostly found in berries, grapes, nuts, coffee, tea, fruits and wine. The second class is condensed tannin or non-hydrolysable tannin. They are flavanoid tanning substances that belong to oligomeric and polymeric proanthocyanidins. Condensed tannins are obtained from cocoa, various fruits like pears, apples, green grapes, legumes such as chickpeas, red kidney beans and peas, respectively (Costain, 2001; Selma et al., 2009). The third class of tannin is complex tannin. Complex tannins or non-classified tannins have catechin units that are glycosidically bound to the gallic acid (gallotannins) or ellagic acid (ellagitannins). Several sources of complex tannins are camelliatannins A and B (*Camellia japonica* L), malabarin A (*Melastoma malabathricum*) and acutissimin A. (Okuda and Ito, 2011).





Fig. 1: Major Classes of Tannins.

Hydrolysable tannins can be helpful in the leather, wine and plastic manufacturing industries (Guo et al., 2020; Motta et al., 2020). They can be used as antimutagenic, anticancer, antioxidant, wood preservative, anti-corrosion agents for metals. (Tomak and Gonultas, 2018; Byrne et al., 2019).

Tannase

Tannase (E.C. 3.1.1.20) or tannin acyl hydrolase (TAH) is an explored group of hydrolases family. The catalytic reaction of tannase involves the hydrolysis of ester bond and depside bond present in various substrates i.e., tannic acid, complex

tannin, gallic acid esters, epigallocatechin gallate, methyl gallate, ethyl gallate, n-propyl gallate to produce gallic acid (3, 4, 5-tri hydroxybenzoic acid) and glucose (Lal and Gardner, 2012). Several intermediate molecules (Figure 2) are formed during the hydrolytic breakdown of tannic acid into gallic acid and glucose by tannase enzyme i.e (II) 1, 2, 3, 4, 6, - pentagalloyl glucose, (III) 2,3,4,6 tetragalloyl glucose and (IV) monogalloyl glucose (Lekha and Lonsane, 1997).



(R1-gallic acid; R2-m-di gallic acid; I-tannic acid, II-1,2,3,4,6-pentagalloyl glucose; III - 2,3,4,6 tetragalloyl glucose and IV- monogalloyl glucose)

Fig. 2: Mechanism of hydrolysis of tannic acid by Tannase (Lekha and Lonsane, 1997)

The novel biotechnological approach of gallic acid is the production of trimethoprim. Gallic acid has emerged as a basic intermediate of trimethoprim containing bacteriostatic and broad-spectrum features. (Anderson et al., 1980; Misro et al., 1997; Mukherjee and Banerjee, 2003). The enormous applications of gallic acid such as in the manufacture of ink, dye and paper, in photographic development, in the tanning process of leather are found in many sectors of various industries.

Tannase Structure

Ren et al. (2013) reported the monomeric three-dimensional structure of tannase from *Lactobacillus plantarum* SICC 1.15 (Figure 3). This tannase contains 50,747 Da molecular weight and 469 amino acid residues. However, the crystal structure of tannase consists of two domains such as α/β -hydrolase domain (residues 4-204 and 396-469) and lid domain (residues 205-395). The α/β -hydrolase domain contains six α helices such as $\alpha 1$ - $\alpha 4$ and $\alpha 12$ - $\alpha 13$ and nine β -stranded sheets such as $\beta 1$ - $\beta 7$ and $\beta 12$ - $\beta 13$. Among the α helices, $\alpha 2$ - $\alpha 4$ and $\alpha 12$ are present on one side of the sheet, while $\alpha 1$ and $\alpha 13$ are present on the opposite side. The other domain known as the lid domain contains seven α -helices such as $\alpha 5$ - $\alpha 11$ and two β -stranded sheets such as $\beta 8$ - $\beta 9$ and $\beta 10$ - $\beta 11$. The two β -stranded sheets lie between the $\beta 7$ - $\beta 12$ strands. The active site of the tannase enzyme is revealed in the α/β -hydrolase domain. A deep tunnel is formed by the juxtaposition of α/β -hydrolase and lid domain. The

tunnel wall is formed by three amino acid residues as Ser163, Asp419 and His451, known as a catalytic triad. Along with the catalytic triad (Ser163, Asp19 and His451), tannase also contains serine residue Ser163 that is located in the pentapeptide motif i.e. GXSG or Gly161-X-Ser163-X-Gly165 (where X = any amino acid residue) between $\beta 6$ and $\alpha 6$. The active site of the tannase consists of three charged residues as Lys 343, Glu357 and Asp421. These three charged residues play a major role in the hydrolysis and binding of galloyl moiety of the substrate.




Fig. 3: Structure of *L. plantarum* tannase. The structure contains α/β -hydrolase domain in red ribbon view and lid domain in blue-ribbon view (Matoba et al., 2013)

Sources of Tannase

Tannase can be found in prokaryotic and eukaryotic cells. It is widely distributed in animals, plants and microorganisms. Among various sources, microbial strains play a crucial role in tannase production because of their great stability over plants and animals. Bacteria, molds and yeasts produce 30%, animals produce 8% and plants produce 4% industrial enzymes.

Plants derived tannase

Tannin-rich plants are great sources for tannase production. Various parts of plants such as bark, woods, stems, roots, fruits, leaves and seeds contain a high amount of tannin. Different plant species like *Acacia katechu*, *Agrimonia eupatoria*, *Diospyros kaki*, *Hamamelis virginiana*, *Monochaetum multiflorum*, *Syzygium cumini*, *Terminalia chebula* and *Quercus robur* were reported to be efficient tannase producers (Shen et al., 2006; Granica et al., 2013; Zhange et al., 2011). Plants produce a variety of acids such as chebulinic acid, gallic acid, and hexahydroxyphenic acid and a large quantity of sugar, during the growth phase. As fruits become mature/ ripen, tannase esterified these acids with sugar (i.e glucose) and form complex tannin (Lekha

and Lonsane, 1997).

Animals and Insects Derived Tannase

The ruminal mucosa of several animals such as koalas (*Phascolarctos cinereus*) (Osawa, 1992), goats (*Capra hircus*) (Brooker et al., 1994; Nelson et al., 1995), and horses (*Equus caballus*) (Nemoto, 1995) contain a small amount of tannase that degrade tannin from the nutritional resources and narrow down harmful effects of tannin on animals (Mosleh et al., 2014). Tannase can be extracted from the larva of some insects such as Indian Mole Cricket (*Gryllotalpa krishnani*), Cynipid, *Salix caprea* (Mani, 1966). The larva of the insects contains other digestive enzymes such as invertase and diastase. The role of digestive enzymes is to act on the large molecules of plant cells and convert them into simple forms and help insects in digestion. Tannin present in the leaves of plants precipitates with digestive enzymes and inactivates them. Tannase from the larva of insects acts on tannin and protects the digestive enzymes (Mani, 1966).

Microbially derived tannase

Many microorganisms such as bacteria, fungi, yeast are known to be tannase producers. Tannase producing microbes utilize tannin as a sole source of carbon and transform tannin into glucose and gallic acid.

Bacterial tannase

For the past few decades, the most significant

approach to obtain tannase is from bacterial source compared to yeast and fungal sources due to its high genetic manipulation, biochemical diversity, stability in an extreme environment, and extracellular enzyme production (Jana et al., 2014, Aguilar and Gutierrez Sanchez, 2001). In earlier reports, many bacterial strains such as *Achromobacter*, *Klebsiella*, *Bacillus*, *Corynebacterium*, *Citrobacter*, *Streptococcus*, *Leuconostoc*, *Pediococcus*, *Pantoea* and *Streptomyces* were able to produce tannase. Out of 21 different bacterial genera, *Lactobacilli* are the most dominant tannase producing bacterial genera among others (Chandrasekaran and Beena, 2013).

Hydrolysable tannins such as gallotannins and ellagitannins are hydrolysed by bacterial tannase (Bhat et al., 1998). Glucose and gallic acid are the hydrolysed products of tannin through tannase. Gallic acid is decarboxylated by the enzyme gallate decarboxylase and converted to pyrogallol. The pyrogallol is converted into pyruvic acid, cis-aconitic acid and 3-hydroxy-5-oxo-hexanoate (Figure 4). These three molecules enter and circulate into the TCA cycle. 3-hydroxy-5-oxo-hexanoate is the breakdown product of dihydro phloroglucinol. Dihydro phloroglucinol and phloroglucinol are produced through catalytic actions of dihydro phloroglucinol hydrolase and phloroglucinol isomerase from pyrogallol.




Fig. 4: Pathway showing Tannin degradation by bacterial tannase (Jana et al., 2014)

Fungal tannase

Fungal tannase have better degrading ability for hydrolysable tannin and condensed tannin (Belmares-Cerda et al., 2003). Out of 20 different genera of fungi, Aspergilli and Penicillia are the most dominant groups of tannase producers (Raghuwanshi, 2012). Among 20 different fungal

genera, 27 species of *Aspergillus*, 24 species of *Penicillium*, 4 species of *Trichoderma* and 3 species of *Fusarium* are known as tannase producers. Several fungal species such as *Mucor* sp., *Paciliomyces* sp., *Hyalopus* sp. and *Neurospora* sp. are also reported as tannase producers. The pathway of tannin degradation by fungal tannase is shown in Figure 5.




Fig. 5: Pathway showing Tannin degradation by fungal tannase (Banerjee and Mahapatra, 2012)

Yeast Tannase

Yeast tannases are less efficient for degrading naturally obtained tannin compare to tannic acid (Deschamps et al., 1983). Nowadays, researchers develop conventional and biotechnological approaches for the production of tannase from yeast due to its short growth period and easy accessibility towards molecular genetics. Various yeast species such as *Candida* sp., *Pichia* sp., *Debaryomyces hansenii*, *Mycotorula japonica* and *Saccharomyces cerevisiae* are reported as tannase producers.

Actinomycetal Tannase

Various actinomycetal species such as *Streptomyces svaceus* (Wang, et al., 2018), *Streptomyces gancidicus* (Tripathi et al., 2021), *Streptomyces* sp. AL1L (Roy et al., 2018) are reported as tannase producers.

Tannase Assay

Diversified analytical methods such as UV-spectrometric (Parmentier, 1973 and Iibuchi et al., 1967), colorimetric (Haslam and Tanner, 1970), photometric (Chen, 1969), titrimetric (Freudenberg et al., 1927; Nishira, 1961; Haslam and Stangroom, 1966), chromatographic (Jean et al., 1981) etc. are reported. All these methods used for assay of tannase are based on either determination of residual tannic acid (Deschamps et al., 1983) or formation of gallic acid, a byproduct of enzymatic hydrolysis of tannic acid.

Solid state and Submerged Fermentative production of tannase

Fermentation is the metabolic process in which complex compounds are converted into simple compounds by using various microorganisms such

as bacteria and fungi. The fermentation process is classified into two broad classes as solid-state fermentation (SSF) and submerged fermentation (SmF) based on the type of substrates used in the process.

Solid-state fermentation or SSF is defined as a process in which microbes utilize natural substrates or non-soluble inert supports as a nutritional source and can grow in the presence of low water availability or absence of water. SSF is a favorable method for tannase production due to its simple operating system, less water consumption and low-cost production (Aguilar et al., 2001 and Barrios González, 2012). Selwal et al. (2010) found maximum tannase activity in the presence of amla (*Phyllanthus emblica*) and keekar (*Acacia nilotica*) leaves by *Pseudomonas aeruginosa* IIB. Ding et al. (2020) used mixed strains of *Bacillus subtilis*, *Aspergillus niger* and *Saccharomyces cerevisiae* for the production of tannase during solid-state fermentation of tea residue.

Submerged fermentation or SmF is defined as a process in which microbes utilize dissolved or suspended nutrients from the liquid medium (Frost, 1987). Among the various fungi, *Penicillium verrucosum* and *Aspergillus niger* degraded tannin from the coffee pulp and tea residue, respectively under SSF and produced tannase (Bhoite and Murthy, 2015 and Sharma et al., 2014). Submerged fermentation is more preferable compared to solid-state fermentation for the production of bacterial and fungal tannase in various industries. *Bacillus megaterium* showed maximum tannase activity (10.77 U/mL/min.) in the presence of 1% tannic acid concentration under SmF (Tripathi et al., 2016).

Tannase Purification

Most of the produced enzymes are extracellular or intracellular through the fermentation process. Cells are directly separated from the fermentation broth for the recovery of the extracellular enzyme, wherein the case of intracellular enzyme cells are disrupted by the mechanical or non mechanical methods. The purification process mostly depends upon the need for the enzyme in the market. Purification of the enzyme is carried out to recover a high yield with the greatest catalytic activity and purity. Downstream processing is essential to remove the impurities from the end product. It is a multistep process in which various methods such as Salting out or solvent precipitation, Ultrafiltration, electrophoresis and chromatography methods could be employed. The initial step for purification

of an enzyme is ammonium sulphate precipitation. The role of ammonium sulphate in purification is to stabilize the protein. Various types of precipitating agents like 2 ethoxy 6-9 diamino acridinium lactate or rivanol, as well as such polymers like dextran, PEG (polyethylene glycol) and polyvinyl alcohol, were used for the precipitation of tannase (Sharma et al., 1999 and Aoki et al., 1976). The second step of enzyme purification is a desalting process or dialysis. Dialysis is a process in which small molecules are removed from a mixture of molecules. The process was performed by using a semipermeable membrane. The third step applicable in tannase purification is ion exchange chromatography. The protein separated by ion-exchange chromatography is based on its charge that interacts with an oppositely charged ion-exchange matrix. Many researchers have used anion exchangers such as DEAE-Sephadex, DEAE-cellulose, DEAE-Sephadex A-50, and DEAE Sepharose for purification of tannase due to its acidic nature (Bhardwaj et al., 2003, Mahendran et al., 2006, Sharma et al., 1999). The fourth or last step of enzyme purification is gel permeation or size exclusion chromatography. The gel beads used in gel filtration chromatography are prepared from Sepharose, Agarose, Vinyl polymers, Polyacrylamide and Sephadex. Sephadex G-100, Sephadex G-200 and Sephadex G-150 were used as a gel in gel filtration chromatography for tannase purification (Beena et al., 2010, Ramirez-Coronel et al., 2003 and Sharma et al., 1999).

Biochemical Properties and Kinetics of Tannase

Different tannases obtained from different microbial sources are characterized and have some biochemical properties that give information related to the nature and structure of the enzyme. Various parameters such as temperature, pH, metal ions, activators, inhibitors and substrate specificity affect the enzyme activity. Microorganisms express the gene for the enzyme at specific pH. The optimum fungal and bacterial tannase biosynthesis has been reported in 3-8 pH range. However, in certain reports, the optimum pH of 8 and 8.9 was reported for bacterial tannase (Belur et al., 2010 and Matsuda et al., 2016). The optimal pH for actinomycetal and yeast tannase was 6 and 4.5, respectively (Roy et al., 2018 and Pan et al., 2020). Tannase from *A. fumigatus* was stable at pH 4.0 and it lost enzyme activity at pH 8.0 (Batra and Saxena, 2005). Mahapatra et al. (2005) reported that mostly fungal tannase required an acidic environment to become active. Many researchers reported the optimum temperature range of some tannase-producing microorganisms

was between 30-70 °C (Raghuwanshi et al., 2011; Beniwal et al., 2013 and Govindarajan et al., 2018). The native structure of tannase might be affected by the presence of metal ions due to an increase in the ionic strength of the solution (Chaitanyakumar and Anbalagan, 2016). At a high concentration of metal ions, tannase activity was decreased. In the previous kinds of literature, tannase activity was enhanced in the presence of sodium chloride (Mondal et al., 2001; Aftab and Hamid, 2016). The catalytic activity of the enzyme is affected by the surfactants (Prasad et al., 2012). Sodium lauryl sulfate (SDS) showed the stimulatory effect on tannase produced from *B. subtilis* at 1% (v/v) concentration, while Tween 80, Tween 60, DMSO (Dimethyl Sulfoxide), Sodium azide, β -mercaptoethanol, EDTA, PMSF (phenylmethylsulfonyl fluoride) and Triton X-100 showed a negative effect on tannase activity (Jana et al., 2013). Various polar and nonpolar organic solvents also affect tannase activity. Several nonpolar solvents such as hexane, toluene and benzene enhanced tannase activity. However, few polar solvents such as methanol, isoamyl alcohol and butanol inhibited tannase activity (Beniwal et al., 2010; Valera et al., 2015).

Applications of Tannase

Tannase is an important industrially relevant and versatile enzyme that has tremendous applications such as in the manufacturing of tea, wine, beer, food, pharmaceutical, chemical, brewing and leather industries. Moreover, the tannase enzyme is also used for the bioremediation of toxic wastes from tannery effluents.

Gallic acid Production

Gallic acid or 3, 4, 5-trihydroxybenzoic acid is an organic compound that is widely used in pharmaceutical and chemical industries. Gallic acid is obtained from various sources such as plants and microorganisms. Gallic acid is present in a plant as free molecules or it can be produced from the tannase-producing microorganisms by hydrolysis of tannic acid or tannin-rich substrates. 3, 4, 5 trimethoxy benzaldehyde is the intermediate component during the manufacturing of trimethoprim from gallic acid. Trimethoprim is effective against gram-negative and gram-positive microorganisms when it synergistic with sulfonamide. Trimethoprim inhibits the dihydrofolate reductase enzyme and prevents the synthesis of folic acid. Folic acid plays a very important role in the synthesis of DNA. Gallic acid serves as an antiapoptotic, antiviral, antimutagenic and anticancer agent (Badhani et al., 2015).

Instant tea Preparation and Tea Cream Solubilization

Tea is the most highly consumed and favorable beverage worldwide, after water. There are so many health benefits of tea such as it helps to reduce the risk of cancer, heart disease and diabetes. When tea leaves are treated with tannase. It improves the flavor of tea with better acid stability and cold water solubility (Natarajan et al., 2009). It also enhances the bright reddish color due to the formation of epigallocatechin gallate from epicatechin and gallic acid. Haze and tea cream formation during the preparation of tea beverages should be removed by tannase treatment.

In Beverages

The application of tannase in beverage industries is to remove water insoluble precipitates and undesirable tannin which is responsible for haze formation and bitter taste (Boadi and Neufeld, 2001). Additionally, tannase plays a major role in the removal of tannins and phenolic compounds from the different fruit juices and influences the flavor and maintains the color stability of the juice during the long period of preservation (Aguilar et al., 2007).

In Animal feed and Cell wall Digestion

The quality of animal feed can be also improved by the digestibility of the plant cell wall. Tannase acts on crosslinks of plant cell wall polymers that contain ferulic acid dehydrodimers. Tannase hydrolyzes diethyl diferulates and improves the quality of animal feed (Garcia-Conesa et al., 2001).

In the Biological Treatment of Tannery Effluent

The tannery industry is one of the largest industries and most important for the tanning of leather in India. The tanning process is necessary to prevent the leather from decomposition and to impart the color of the leather. Moreover, it causes environmental pollution during the discharge of tannery wastes to sewage or natural bodies. Besides these, tannin present in the effluent also inhibits the microorganisms. Various chemical and biological methods are used to remove tannin and other phenolic compounds from the tannery effluent. Recently, biodegradation technology is widely used for the treatment of tannery effluent and other types of industrial wastes. Tannase treatment is an eco-friendly treatment that helps to remove tannin and decrease pollution load such as BOD and COD from industrial wastes or wastewater (Zhao et al., 2017).

References

- Aftab, M.N. and Hamid, M., 2016. Production and characterization of tannase from a newly isolated *Bacillus subtilis*. *Pakistan Journal of Botany*, 48(3), 1263-1271.
- Aguilar, C.N. and Gutiérrez-Sánchez, G., 2001. Sources, properties, applications and potential uses of tannin acyl hydrolase. *Food Science and Technology International*, 7(5), 373-382. <https://doi.org/10.1106%2F69M3-B30K-CF7Q-RJ5G>
- Aguilar, C.N., Rodríguez, R., Gutiérrez-Sánchez, G., Augur, C., Favela-Torres, E., Prado-Barragan, L.A., Ramírez-Coronel, A. and Contreras-Esquivel, J.C., 2007. Microbial tannases: advances and perspectives. *Applied microbiology and biotechnology*, 76(1), 47-59. <https://doi.org/10.1007/s00253-007-1000-2>
- Anderson, R., Grabow, G.I.S.E.L.A., Oosthuizen, R.O.B.Y.N., Theron, A.N.N.E.T.T.E. and Van Rensburg, A.J., 1980. Effects of sulfamethoxazole and trimethoprim on human neutrophil and lymphocyte functions in vitro: in vivo effects of co-trimoxazole. *Antimicrobial agents and chemotherapy*, 17(3), 322-326. <https://doi.org/10.1128/aac.17.3.322>
- Aoki, K., Shinke, R. and Nishira, H., 1976. Purification and some properties of yeast tannase. *Agricultural and Biological Chemistry*, 40(1), 79-85. <https://doi.org/10.1080/00021369.1976.10861997>
- Badhani, B., Sharma, N. and Kakkar, R., 2015. Gallic acid: a versatile antioxidant with promising therapeutic and industrial applications. *Rsc Advances*, 5(35), 27540-27557. <https://doi.org/10.1039/C5RA01911G>
- Banerjee, D. and Mahapatra, S., 2012. Fungal tannase: a journey from strain isolation to enzyme applications. *Dynamic Biochemistry, Process Biotechnology and Molecular Biology*, 6(2), 49-60.
- Barbehenn, R.V. and Constabel, C.P., 2011. Tannins in plant-herbivore interactions. *Phytochemistry*, 72(13), 1551-1565. <https://doi.org/10.1016/j.phytochem.2011.01.040>
- Barrios-González, J., 2012. Solid-state fermentation: physiology of solid medium, its molecular basis and applications. *Process Biochemistry*, 47(2), 175-185. <https://doi.org/10.1016/j.procbio.2011.11.016>
- Batra, A. and Saxena, R.K., 2005. Potential tannase producers from the genera *Aspergillus* and *Penicillium*. *Process Biochemistry*, 40(5), 1553-1557. <https://doi.org/10.1016/j.procbio.2004.03.003>
- Beena, P., 2010. Production, purification, genetic characterization and application studies of tannase enzyme from marine fungus *Aspergillus awamori*. From: Ph. D. Thesis. Department of Biotechnology, Cochin University of Science and Technology, Cochin, Kerala, India.
- Belmares-Cerda, R., Reyes-Veja, M. L., Contreras Esquivel, J. C., Rodríguez-Herrera, R., and Aguilar, C. N., 2003. Effect of carbon source on tannase production by two strains of *Aspergillus niger*. *Revista Mexicana de Ingeniería Química* 2, 95-100.
- Belur, P.D., Mugeraya, G. and Kuppalu, N.R., 2010. Temperature and pH stability of a novel cell-associated tannase of *Serratia ficaria* DTC. *International Journal of Biotechnology and Biochemistry*, 6(5), 667-674.
- Beniwal, V., Chhokar, V., Singh, N. and Sharma, J., 2010. Optimization of process parameters for the production of tannase and gallic acid by *Enterobacter cloacae* MTCC 9125. *Journal of American Science*, 6(8), 389-397.
- Beniwal, V., Kumar, A., Goel, G. and Chhokar, V., 2013. A novel low molecular weight acidothermophilic tannase from *Enterobacter cloacae* MTCC 9125. *Biocatalysis and Agricultural Biotechnology*, 2(2), 132-137. <https://doi.org/10.1016/j.bcab.2013.03.002>
- Bhardwaj, R., Singh, B. and Bhat, T.K., 2003. Purification and characterization of tannin acyl hydrolase from *Aspergillus niger* MTCC 2425. *Journal of Basic Microbiology: An International Journal on Biochemistry, Physiology, Genetics, Morphology, and Ecology of Microorganisms*, 43(6), 449-461. <https://doi.org/10.1002/jobm.200310273>
- Bhat, T.K., Singh, B. and Sharma, O.P., 1998. Microbial degradation of tannins—a current perspective. *Biodegradation*, 9(5), 343-357. <https://doi.org/10.1023/A:1008397506963>
- Bhoite, R.N. and Murthy, P.S., 2015. Biodegradation of coffee pulp tannin by *Penicillium verrucosum* for production of tannase, statistical optimization and its application. *Food and Bioprocess Processing*, 94, 727-735. <https://doi.org/10.1016/j.fbp.2014.10.007>
- Boadi, D.K. and Neufeld, R.J., 2001. Encapsulation of tannase for the hydrolysis of tea tannins. *Enzyme and Microbial Technology*, 28(7-8), 590-595. [https://doi.org/10.1016/S0141-0229\(01\)00295-2](https://doi.org/10.1016/S0141-0229(01)00295-2)
- Brooker, J.D., O'donovan, L.A., Skene, I., Clarke, K., Blackall, L. and Muslera, P., 1994. *Streptococcus caprinus* sp. nov., a tannin resistant ruminal bacterium from feral goats. *Letters in Applied Microbiology*, 18(6), 313-318. <https://doi.org/10.1111/j.1472-765X.1994.tb00877.x>
- Byrne, C., D'Alessandro, O., Selmi, G.J., Romagnoli, R. and Deyá, C., 2019. Primers based on tara and quebracho tannins for poorly prepared steel surfaces. *Progress in Organic Coatings*, 130, 244-250. <https://doi.org/10.1016/j.porgcoat.2019.02.003>
- Chaitanyakumar, A. and Anbalagan, M., 2016. Expression, purification and immobilization of tannase from *Staphylococcus lugdunensis* MTCC 3614. *AMB Express*, 6(1), 1-9. <https://doi.org/10.1186/s13568-016-0261-5>
- Chandrasekaran, M. and Beena, P.S., 2013. Tannase: source, biocatalytic characteristics, and bioprocesses for production. In *Marine enzymes for biocatalysis*,

- Woodhead Publishing.,259-293. <https://doi.org/10.1533/9781908818355.3.259>
24. Chen, T., 1969. The induction of tannase in *Aspergillus niger*. In Ph.D. Thesis in Biochemistry. Univ. of California Berkeley.
 25. Costain, L., 2001. *Supernutrients Handbook*. Dorling Kindersley, New York, NY.
 26. Dai, J. and Mumper, R.J., 2010. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules*, 15(10),7313-7352. <https://doi.org/10.3390/molecules15107313>
 27. Deschamps, A.M., Otuk, G. and Lebeault, J.M., 1983. Production of tannase and degradation of chestnut tannin by bacteria. *Journal of Fermentation Technology*, 61(1), 55-59.
 28. Deschamps, A.M., Otuk, G. and Lebeault, J.M., 1983. Production of tannase and degradation of chestnut tannin by bacteria. *Journal of Fermentation Technology*, 61(1), 55-59.
 29. Ding, X., Yao, L., Hou, Y., Hou, Y., Wang, G., Fan, J. and Qian, L., 2020. Optimization of culture conditions during the solid-state fermentation of tea residue using mixed strains. *Waste and Biomass Valorization*, 1-9. <https://doi.org/10.1007/s12649-019-00930-4>
 30. Działo, M., Mierziak, J., Korzun, U., Preisner, M., Szopa, J. and Kulma, A., 2016. The potential of plant phenolics in prevention and therapy of skin disorders. *International journal of molecular sciences*, 17(2), 160. <https://doi.org/10.3390/ijms17020160>
 31. Freudenberg, Karl; Blümmel, Franz; Frank, Theodor, 1927. Knowing the tannase II. *Hoppe-Seyler's magazine for physiological chemistry*, 164 (4-6),262-270. <https://doi.org/10.1515/bchm2.1927.164.4-6.262>
 32. Frost, M.G., 1987. Production of enzymes by fermentation. *Biotechnology*, 108.
 33. Garcı a-Conesa, M.T., Østergaard, P., Kauppinen, S. and Williamson, G., 2001. Hydrolysis of diethyl diferulates by a tannase from *Aspergillus oryzae*. *Carbohydrate Polymers*, 44(4), 319-324. [https://doi.org/10.1016/S0144-8617\(00\)00248-4](https://doi.org/10.1016/S0144-8617(00)00248-4)
 34. Gou, X., Okejiri, F., Zhang, Z., Liu, M., Liu, J., Chen, H., Chen, K., Lu, X., Ouyang, P. and Fu, J., 2020. Tannin-derived bimetallic CuCo/C catalysts for an efficient in-situ hydrogenation of lauric acid in methanol-water media. *Fuel Processing Technology*, 205, 106426. <https://doi.org/10.1016/j.fuproc.2020.106426>
 35. Govindarajan, R.K., Krishnamurthy, M., Neelamegam, R., Shyu, D.J., Muthukalingan, K. and Nagarajan, K., 2018. Purification, structural characterization and biotechnological potential of tannase enzyme produced by *Enterobacter cloacae* strain 41. *Process Biochemistry*, 77, 37-47. <https://doi.org/10.1016/j.procbio.2018.10.013>
 36. Granica, S., Krupa, K., Kłębowska, A. and Kiss, A.K., 2013. Development and validation of HPLC-DAD-CAD-MS3 method for qualitative and quantitative standardization of polyphenols in *Agrimoniae eupatoria* herba (Ph. Eur). *Journal of Pharmaceutical and Biomedical Analysis*, 86,112-122. <https://doi.org/10.1016/j.jpba.2013.08.006>
 37. Haslam, E. and Stangroom, J.E., 1966. The esterase and depsidase activities of tannase. *Biochemical Journal*, 99(1), 28-31. <https://doi.org/10.1042/bj0990028>
 38. Haslam, E. and Tanner, R.J.N., 1970. Spectrophotometric assay of tannase. *Phytochemistry*, 9(11), 2305-2309. [https://doi.org/10.1016/S0031-9422\(00\)85729-5](https://doi.org/10.1016/S0031-9422(00)85729-5)
 39. Iibuchi, S., Minoda, Y. and Yamada, K., 1967. Studies on tannin acyl hydrolase of microorganisms: Part II. A new method determining the enzyme activity using the change of ultra violet absorption. *Agricultural and Biological Chemistry*, 31(5),513-518. <https://doi.org/10.1080/00021369.1967.10858840>
 40. Jana, A., Halder, S.K., Banerjee, A., Paul, T., Pati, B.R., Mondal, K.C. and Mohapatra, P.K.D., 2014. Biosynthesis, structural architecture and biotechnological potential of bacterial tannase: a molecular advancement. *Bioresource Technology*, 157, 327-340. <https://doi.org/10.1016/j.biortech.2014.02.017>
 41. Jana, A., Maity, C., Halder, S.K., Das, A., Pati, B.R., Mondal, K.C. and Mohapatra, P.K.D., 2013. Structural characterization of thermostable, solvent tolerant, cytosafe tannase from *Bacillus subtilis* PAB2. *Biochemical engineering journal*, 77,161-170. <https://doi.org/10.1016/j.bej.2013.06.002>
 42. Jean, D., Pourrat, H., Pourrat, A. and Carnat, A., 1981. Assay of tannase (tanin acylhydrolase EC 3.1. 1.20) by gas chromatography. *Analytical Biochemistry*, 110(2), 369-372. [https://doi.org/10.1016/0003-2697\(81\)90205-0](https://doi.org/10.1016/0003-2697(81)90205-0)
 43. Khanbabaee, K. and Van Ree, T., 2001. Tannins: classification and definition. *Natural product reports*, 18(6),641-649. <https://doi.org/10.1039/b101061l>
 44. Lal, D. and Gardner, J.J., 2012. Production, characterization and purification of tannase from *Aspergillus niger*. *European Journal of Experimental Biology*, 2(5), 1430-1438.
 45. Lekha, P.K. and Lonsane, B.K., 1997. Production and application of tannin acyl hydrolase: state of the art. *Advances in applied microbiology*, 44,216-260. [https://doi.org/10.1016/s0065-2164\(08\)70463-5](https://doi.org/10.1016/s0065-2164(08)70463-5)
 46. Mahapatra, K., Nanda, R.K., Bag, S.S., Banerjee, R., Pandey, A. and Szakacs, G., 2005. Purification, characterization and some studies on secondary structure of tannase from *Aspergillus awamori* nakazawa. *Process Biochemistry*, 40(10),3251-3254. <https://doi.org/10.1016/j.procbio.2005.03.034>
 47. Mahendran, B., Raman, N. and Kim, D.J., 2006. Purification and characterization of tannase from *Paecilomyces variotii*: hydrolysis of tannic acid using immobilized tannase. *Applied Microbiology*

- and Biotechnology, 70(4),444-450. <https://doi.org/10.1007/s00253-005-0082-y>
48. Mani, M.S., 1966. Ecology of plant galls (Monographiae Biologicae).Journal of Ecology, 54(1), 276-277. <https://doi.org/10.2307/2257675>
 49. Matoba, Y., Tanaka, N., Noda, M., Higashikawa, F., Kumagai, T. and Sugiyama, M., 2013. Crystallographic and mutational analyses of tannase from *Lactobacillus plantarum*. Proteins: Structure, Function, and Bioinformatics, 81(11), 2052-2058. <https://doi.org/10.1002/prot.24355>
 50. Matsuda, M., Hirose, Y., Kanauchi, M., Hatanaka, S. and Totsuka, A., 2016. Purification and characteristics of tannase produced by lactic acid bacteria, *Lactobacillus plantarum* H78. Journal of the American Society of Brewing Chemists, 74(4),258-266.<https://doi.org/10.1094/ASBCJ-2016-4298-01>
 51. Misro, S.K., Kumar, M.R., Banerjee, R. and Bhattacharyya, B.C., 1997. Production of gallic acid by immobilization of *Rhizopus oryzae*. Bioprocess Engineering, 16(5),257-260. <https://doi.org/10.1007/s004490050318>
 52. Mondal, K.C., Banerjee, D., Banerjee, R. and Pati, B.R., 2001. Production and characterization of tannase from *Bacillus cereus* KBR9. The Journal of General and Applied Microbiology, 47(5), 263-267. <https://doi.org/10.2323/jgam.47.263>
 53. Mosleh, H., Naghiha, A., NAGHI, K.A. and Khajavi, M., 2014. Isolation and identification of tannin-degrading bacteria from native sheep and goat feces in Kohgiluyeh and Boyer-Ahmad Province. International Journal of Advanced Biological and Biomedical Research,,2(1), .176-180.
 54. Motta, S., Guaita, M., Cassino, C. and Bosso, A., 2020. Relationship between polyphenolic content, antioxidant properties and oxygen consumption rate of different tannins in a model wine solution. Food chemistry, 313, 126045.<https://doi.org/10.1016/j.foodchem.2019.126045>
 55. Mukherjee, G. and Banerjee, R., 2003. Production of gallic acid. Biotechnological routes (Part 2). Chimica oggi, 21(3-4), 70-73.
 56. Natrajan, K., 2009. Tannase: A tool for instantaneous tea. Current Biotica, 3(1), 96-103.
 57. Nelson, K.E., Pell, A.N., Schofield, P. and Zinder, S., 1995. Isolation and characterization of an anaerobic ruminal bacterium capable of degrading hydrolyzable tannins. Applied and Environmental Microbiology, 61(9), 3293-3298. <https://doi.org/10.1128/aem.61.9.3293-3298.1995>
 58. Nemoto, K., Osawa, R.O., Hirota, K., Ono, T. and Miyake, Y., 1995. An investigation of gram-negative tannin-protein complex degrading bacteria in fecal flora of various mammals. Journal of Veterinary Medical Science, 57(5), 921-926.<https://doi.org/10.1292/jvms.57.921>.
 59. Nishira, H., 1961. Degradative metabolism of tannin by molds. Hakko Kogaku Zasshi, 39, 137-146.
 60. Okuda, T. and Ito, H., 2011. Tannins of constant structure in medicinal and food plants—hydrolyzable tannins and polyphenols related totannins. Molecules, 16(3), 2191-2217.<https://doi.org/10.3390/molecules16032191>
 61. Pan, J., Wang, N.N., Yin, X.J., Liang, X.L. and Wang, Z.P., 2020. Characterization of a Robust and pH-Stable Tannase from Mangrove-Derived Yeast *Rhodospiridium diobovatum* Q95. Marine Drugs, 18(11), 546. <https://doi.org/10.3390/md18110546>
 62. Parmentier, F., 1973. A direct spectrometric method for the assay of tannase. Communication Brève. Archives Internationales de Physiologie et de Biochimie, 81(2), 339-341. <https://doi.org/10.3109/13813457009075189>
 63. Prasad, D., Gupta, R.K., Mathangi, G., Kamini, N.R. and Gowthaman, M.K., 2012. Advances in production and characteristic features of microbial tannases: an overview. Current Trends in Biotechnology and Pharmacy, 6(2), 145-165.
 64. Raghuwanshi, S., Dutt, K., Gupta, P., Misra, S. and Saxena, R.K., 2011. *Bacillus sphaericus*: The highest bacterial tannase producer with potential for gallic acid synthesis. Journal of Bioscience and Bioengineering, 111(6),635-640.<https://doi.org/10.1016/j.jbiosc.2011.02.008>
 65. Raghuwanshi, S., 2012. Tannase from *Penicillium charlesii*: Process optimization, purification, characterization, scale up and industrial applications. From: Ph. D. Thesis. Department of Microbiology, University of Delhi South Campus, New Delhi, India.
 66. Ramírez-Coronel, M.A., Viniegra-Gonzalez, G., Darvill, A. and Augur, C., 2003. A novel tannase from *Aspergillus niger* with β -glucosidase activity. Microbiology, 149(10), 2941-2946.<https://doi.org/10.1099/mic.0.26346-0>
 67. Ren, B., Wu, M., Wang, Q., Peng, X., Wen, H., McKinstry, W.J. and Chen, Q., 2013. Crystal structure of tannase from *Lactobacillus plantarum*. Journal of Molecular Biology, 425(15), 2737-2751. <https://doi.org/10.1016/j.jmb.2013.04.032>
 68. Roy, S., Parvin, R., Ghosh, S., Bhattacharya, S., Maity, S. and Banerjee, D., 2018. Occurrence of a novel tannase (tan B LP) in endophytic *Streptomyces* sp. AL1L from the leaf of *Ailanthus excelsa* Roxb. 3 Biotech, 8(1), 33.<https://doi.org/10.1007/s13205-017-1055-4>
 69. Selma, M.V., Espin, J.C. and Tomas-Barberan, F.A., 2009. Interaction between phenolics and gut microbiota: role in human health. Journal of Agricultural and Food Chemistry, 57(15), 6485-6501.<https://doi.org/10.1021/jf902107d>
 70. Selwal, M.K., Yadav, A., Selwal, K.K., Aggarwal, N.K., Gupta, R. and Gautam, S.K., 2010. Optimization of cultural conditions for tannase production by *Pseudomonas aeruginosa* IIB 8914

- under submerged fermentation. *World Journal of Microbiology and Biotechnology*, 26(4), 599-605. <https://doi.org/10.1007/s11274-009-0209-x>
71. Sharma, K.P., 2019. Tannin degradation by phytopathogen's tannase: A Plant's defense perspective. *Biocatalysis and Agricultural Biotechnology*, 21, 101342. <https://doi.org/10.1016/j.bcab.2019.1013>
 72. Sharma, N.K., Beniwal, V., Kumar, N., Kumar, S., Pathera, A.K. and Ray, A., 2014. Production of tannase under solid-state fermentation and its application in detannification of guava juice. *Preparative Biochemistry and Biotechnology*, 44(3), 281-290. <https://doi.org/10.1080/10826068.2013.812566>
 73. Sharma, S., Bhat, T.K. and Dawra, R.K., 1999. Isolation, purification and properties of tannase from *Aspergillus niger* van Tieghem. *World Journal of Microbiology and Biotechnology*, 15(6), 673-677. <https://doi.org/10.1023/A:1008939816281>
 74. Shen, D., Wu, Q., Wang, M., Yang, Y., Lavoie, E.J. and Simon, J.E., 2006. Determination of the predominant catechins in *Acacia catechu* by liquid chromatography/electrospray ionization-mass spectrometry. *Journal of Agricultural and Food Chemistry*, 54(9), 3219-3224. <https://doi.org/10.1021/jf0531499>
 75. Tomak, E.D. and Gonultas, O., 2018. The wood preservative potentials of valonia, chestnut, tara and sulphited oak tannins. *Journal of Wood Chemistry and Technology*, 38(3), 183197. <https://doi.org/10.1080/02773813.2017.1418379>
 76. Tripathi A. D. and Lakshmi B. 2021. Study on a novel actinomycete, *Streptomyces gancidicus* (MF464014) and enhancing its tannase producing ability by optimizing growth conditions in submerged fermentation. *International Journal of Pharmaceutical Sciences and Research*, 12(1), 640-46. [https://doi.org/10.13040/IJPSR.0975-8232.12\(1\).640-46](https://doi.org/10.13040/IJPSR.0975-8232.12(1).640-46)
 77. Tripathi, A.D., Sharma, K. and B. Lakshmi., 2016. Study on tannase producing *Bacillus megaterium* isolated from tannery effluent. *International Journal of Advanced Research in Biological Sciences*, 3(7), 28-35.
 78. Valera, L.S., Jorge, J.A. and Guimarães, L.H.S., 2015. Characterization of a multi-tolerant tannin acyl hydrolase II from *Aspergillus carbonarius* produced under solid-state fermentation. *Electronic Journal of Biotechnology*, 18(6), 464-470. <https://doi.org/10.1016/j.ejbt.2015.09.008>
 79. Wang, D., Liu, Y., Lv, D., Hu, X., Zhong, Q., Zhao, Y. and Wu, M., 2018. RETRACTED ARTICLE: Substrates specificity of tannase from *Streptomyces svaceus* and *Lactobacillus plantarum*. *AMB Express*, 8(1), 1-8. <https://doi.org/10.1186/s13568-018-0677-1>
 80. Zhang, T., Li, G., Zhi, C. and Mo, H., 2011. Persimmon tannin composition and function. In *International Conference on Agricultural and Biosystems Engineering*. Advanced Biomedical Engineering. 1-2:389-392.
 81. Zhao, C., Shu, L., Wang, G., Yang, Q. Enzymatic Treatment of Organics in Tannery Effluent. *Journal of Residuals Science and Technology*. 2017;14(1):387-391.

